

Mechanisms of aggression and production in chickens: genetic variations in the functions of serotonin, catecholamine, and corticosterone

H.W. CHENG^{1*} and W.M. MUIR²

¹Livestock Behaviour Research Unit, USDA-ARS, W. Lafayette, IN 47907; ²Animal Sciences Department, Purdue University, W. Lafayette, IN 47907, USA

*Corresponding author: hwcheng@purdue.edu

The neuroendocrine systems, such as dopamine (DA) and serotonin (5-HT) as well as corticosterone (CORT), are involved in regulating behavioural patterns and reproduction in humans and other mammals. Similar functions of neuroendocrine system may present in laying hens. To test the hypothesis, two divergent chicken lines were used in the study. Each line has distinct levels of aggressiveness and productivity at a group setting and exhibits different susceptibility to various environmental stressors. We found that, at 21 wks of age, LGPS (Low Group Productivity and Survivability) birds had significantly higher blood concentrations of DA and epinephrine than the KGB birds (Kind Gentle Birds, also previously termed HGPS, birds with a High Group Productivity and Survivability) ($P < 0.01$, respectively). The blood concentration of norepinephrine was not significantly different between the lines but the ratio of epinephrine to norepinephrine was higher in LGPS birds ($P < 0.01$). The blood concentration of 5-HT was also significantly higher in LGPS birds compared to KGB birds ($P < 0.01$). In contrast, KGB birds tended to have a higher level of blood CORT ($P = 0.1$). The results suggest that genetic selection for productivity and survivability with domestic behaviours alters the birds' neuroendocrine homeostasis. The selection-associated plasticity of the neuroendocrine system in controlling animal aggression and productivity were discussed in the article.

Keywords: gene; production; aggression; neuroendocrinology; chickens

Abbreviation Key: CNS = Central nervous system; CORT = Corticosterone; DA = Dopamine; D β H = Dopamine- β -hydroxylase; EP = epinephrine; DXL = Commercial Dekalb XL line; HA line = chicken line selected for high antibody response to SRBC; HAP axis = Hypothalamic-pituitary-adrenal axis; H:L = Heterophil:lymphocyte ratio; KGB = Kinder Gentler Bird selected for high group productivity and longevity; LA line = chicken line selected for high antibody response to SRBC; LH = luteinising hormone;

© World's Poultry Science Association 2007

World's Poultry Science Journal, Vol. 63, June 2007

Received for publication August 14, 2006

Accepted for publication January 24, 2007

LHRH = luteinising hormone-releasing hormone; LGPS = chicken line selected for low group productivity and survivability; MHC = Major histocompatibility complex; NE = norepinephrine; RIA = radioimmunoassay; SRBC = Sheep red blood cell; TH = tyrosine

Introduction

Aggression, in nature, is associated with competition (natural selection) for survival and reproduction (Fraser and Rushen, 1987; Duncan, 1998). However, in artificial production environments, such as in poultry production, aggression causes increased social stress, feather and body damage, in some instances these injuries leading ultimately to cannibalism. Cannibalism is a major concern related non-beaktrimmed bird deaths in current housing environments (Appleby and Hogarth, 1991; Tablante *et al.*, 2000; Glatz, 2005; Rodenburg *et al.*, 2005). To prevent the expression of these injurious behaviours, it is necessary to understand the cellular mechanisms involved in regulating bird domestic behaviour and reproduction.

In humans and other mammals, alterations of neuroendocrine homeostasis have been thought to be a final common pathway in controlling individuals' behavioural patterns and productivity. Specifically, monoamine abnormalities are found to be associated with aggressive behaviours (Berman and Coccaro, 1998) and alternative reproduction (Sharp *et al.*, 1984; Barraclough 1992). Serotonin (5-HT) and its metabolites, 5-hydroxyindoleacetic acid (5-HIAA) as well as density of its receptors have been used as major indicators of abnormal behaviours, including aggression (Valzelli, 1984; Bell and Hobson, 1994; Popova *et al.*, 1997; Barak and Mashiah, 2003; Abumariam *et al.*, 2006). In addition, 5-HT directly and indirectly affects reproductive performances (Tuomisto and Mannisto, 1985; Sirotkin and Schaeffer, 1997). Dopamine (DA) is another monoamine involved in controlling domestic behaviour and reproduction (Snider and Kuchel, 1983; Kudriavtseva *et al.*, 1988; Miczek *et al.*, 1994; Kuikka *et al.*, 1998), and dopaminergic receptor 2 (D₂) gene represents one of the few single-site loci associated with abnormal behaviours in human (Blum and Sheridan, 1995a, b). Catecholamines (like epinephrine, EP and norepinephrine, NE) and corticosterone (CORT), known as "stress hormones", are functionally involved in controlling domestic behaviours (Servo and Naumenko, 1990), linked to mood control (fight or flight), and engaged in cope-with-stress as a sense of well-being (Kopin, 1984; Rolih and Ober, 1995; Kilgour and Szantar-Coddington, 1997; Pollard, 1997). Biological dysregulation of these biogenic amines or hormones may cause brain structural as well as functional reorganization (de Kloet *et al.*, 1996; Ferris, 2000), which in turn affect animal's cope styles and well-being.

The present study was designed to determine if bird neuroendocrine system has similar functions in controlling aggression and production since there is evidence that suggests bird brains possess a core "social behavioural network" which is homologous to the social behavioural network of mammals (Goodson, 2005). In addition, there are similar distributions of neurotransmitter receptors, including dopaminergic and serotonergic receptors between birds and mammals (Dietl and Palacios, 1988; Richfield *et al.*, 1987; Walker *et al.*, 1991).

Two divergent chicken lines, KGB (kind gentle bird) and LGPS (low group productivity and survivability bird, previously termed MBB, mean bad bird), were used in the study. Each line has distinct levels of aggressiveness and productivity at a group setting and exhibits different susceptibility to environmental stressors (Hester *et al.*, 1996 a,b; Cheng *et al.*, 2001a,b; Cheng and Muir, 2005). The differences in behavioural and physiological characteristics of these lines may be reflected in changes in the capability of the neuronal system in response to stimuli. The objective of the study was to determine whether the

effects of selection on behavioural adaptation and productivity could lead to selection-associated functional alterations of the neuroendocrine systems in releasing endogenous psychotropic compounds, such as 5-HT, DA, and CORT; and to evaluate whether the changes of these monoamines and CORT can be used as physiological indicators of an animal's coping ability and well-being.

Material and methods

DEVELOPMENT OF THE GENETIC LINES

The process for developing the genetic lines, KGB and LGPS, has been described previously (Muir, 1996; Cheng and Muir, 2005). Briefly, a programme termed "group selection" was used for developing the research lines. The programme turns "survival of the fittest" emphasized on the individuals to "survival of the adequate" in the group. Results from group selection studies have showed that productivity can be increased while at the same time well-being improved.

The 8th generation was produced from mating 1248 hens with 312 roosters selected at random from 20 wk old pullets and roosters of the 7th generation. Eggs were collected for 2 weeks. Chicks were reared using standard management practices (Muir, 1996), and at 18 wks of age, 9216 pullets were housed by half-sib family in 768 12-birds cages. After 52 wks of production (72 wk of age), birds from 12 cages with the highest group productivity (egg number) and the lowest mortality from cannibalism and flightiness, along with their full and half sib brothers, were selected for the KGB line. To establish a comparison line, differing only in the criteria of selection, birds from 12 cages with the lowest group productivity and the highest mortality, along with their full and half sib brothers, were used to establish a reverse selected LGPS line.

Birds were randomly mated within each line, 4 hens/one rooster; avoiding full or half-sib mating to establish the 9th generation of the KGB and the 2nd generation of the LGPS lines which were used as the genetic material for this research. The detail of the selection technology and rearing programme was reported previously (Muir, 1996; Cheng and Muir, 2005). Pullets of each genetic line were not beak trimmed, and reared under the same conditions, *i.e.*, hatched, vaccinated for *Marek's* and *Newcastle's* disease, and maintained using standard management practices in raised wire cages up to 17 wk of age. At 17 wk of age, birds from each genetic line were randomly assigned to individual cages, each providing 1085 cm² per bird. Feed and water were provided for *ad libitum* consumption. Overhead lights were on daily from 0700 until 1900 h initially, and were increased by 15 min/wk. Light duration was at 13 h daily when the study was performed.

Chicken care guidelines were in strict accordance with the rules and regulations set by Federation of Animal Science Societies (Graig *et al.*, 1999). Experimental protocols were approved by the institutional Animal Care and Use Committee at Purdue University. Efforts were made to minimize animal suffering and the number of birds to be used.

BLOOD SAMPLING

Based on previous observations, the main behavioural adaptation of the KGB birds in responses to social stress became stable after the first 3 wks in responses to social stress (Craig and Muir, 1996a, b). At 21 wks of age, 24 birds from the KGB and LGPS lines, without a plumped egg confirmed by palpating, were bled between 09:00 and 10:00 h (Savory and Mann, 1997). A 5 ml blood sample was collected into a heparinised tube from the brachial vein of each bird within 2 min of removing from its cage. For measuring concentration of 5-HT, the whole blood was used. For measuring concentrations of catecholamines and CORT, blood samples were centrifuged at 700 g for 15 min at 20°C

using Sorvall BC 3B plus centrifuge. Plasmas were kept on ice for further processing or kept at -80°C until measurement.

HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY ASSAY (HPLC ASSAY)

To measure blood concentrations of catecholamines, *i.e.*, DA, EP and NE, the ESA plasma catecholamine analysis kit was used. Briefly, duplicate plasma samples were acidified with 4M perchloric acid, followed by deproteinization with the supplied reagent. After centrifugation, the acid-supernatants with internal standard were added and absorbed onto an alumina minicolumn to bind the catecholamines. HPLC columns were then rinsed and eluted with the supplied solutions, respectively. Eluents were injected into the reverse-phase columns and catechols were detected by liquid chromatography with electrochemical detection. The mobile phase (75 mM Na_2HPO_4 , 1.7 mM OSA, 25 μM EDTA, 10% CH_3CN and 100uL/L Triethylamine, adjusted to pH 3.00 with Phosphoric acid) flow rate was 1.3 mL/min. Catecholamine concentrations were calculated from a reference curve constructed using the provided standards. Concentrations were obtained as picograms per mL.

To measure blood concentration of 5-HT, whole blood samples were acidified in duplicate using 4M perchloric acid and freshly prepared 3% ascorbic acid. After centrifugation, the acid-supernatants were injected into the columns. The mobile phase flowed rate was 1.0 mL/min., and the concentration of 5-HT was calculated from a reference curve made using standard 5-HT.

RADIOIMMUNOASSAY

Total plasma CORT was measured in triplicate using a commercial ^{125}I -soluble antibody CORT radioimmunoassay (RIA) kit with modification as company suggestion for using in birds and as previously described (Renden *et al.*, 1994; Kannan, 1997). In order to validate for parallelism and recovery in birds, adjustments of dilution to 1 to 5, were made, *i.e.*, 20 ul sample to 80 ul steroid diluent. The concentration of CORT was calculated from a reference curve that ranges from 0.1 ng/mL (83% binding) to 4.0 ng/mL (14.9% binding). Addition of known amounts of unlabeled CORT to plasma samples, *i.e.*, producing theoretical concentrations of 0.5, 1.0 and 2.0 ng/mL, resulted in 0.48, 1.08, and 1.97 ng/mL, respectively (CV = 5.0%). The sensitivity of the assay was 0.02 ng/mL. All samples within the experiment were performed at same time.

STATISTICAL ANALYSIS

The experimental design was completely randomized with genetic lines as the main effect. Results were assessed statistically using one-way ANOVA, and intergroup differences were analyzed by Newman-Keuls post hoc test.

Results

LINE DIFFERENCES IN PRODUCTIVITY AND SURVIVABILITY

Productivity and survivability were significantly different among the 8th generation KGB and LGPS lines as well as the overall line from which the selected lines came. The KGB birds had significantly higher productivity than the birds of the overall line which was in turn greater than the LGPS birds, in terms of egg number per cage housed, and egg mass per day per cage, but not egg weight ($P < 0.01$, Table 1). In addition, mortality from cannibalism was significantly higher and survivability was greatly lower in the LGPS line compared to both KGB line and overall line ($P < 0.01$, Table 1).

GENETIC SELECTION-INDUCED ALTERATIONS IN PLASMA CONCENTRATIONS OF SEROTONIN

Blood concentration of 5-HT were measured using HPLC (*Figure 1*), and the reference curves created using commercially purified 5-HT in this study was reliable and repeatable in duplicate running the samples ($R^2 = 1.0$, *Figure 1a*). There was a significantly higher blood concentration of 5-HT in the LGPS birds than in the KGB birds (1.43 vs. 1.18 mg/ml, $P < 0.01$) (*Figure 1b*).

GENETIC SELECTION-INDUCED ALTERATIONS IN PLASMA CONCENTRATIONS OF CATECHOLAMINES

Genetic selection-induced different alterations of blood concentrations of catecholamines between the lines were shown in the *Figure 2* and *Table 2*. Based on the reference curve developed using commercial standard ($R^2 = 1.0$, *Figure 2a*), the LGPS birds exhibited higher concentrations of DA and EP as compared to the KGB birds, *i.e.*, 2.42 vs. 0.59 ng/mL ($P < 0.01$) and 0.59 vs. 0.30 ng/mL ($P < 0.01$), respectively (*Table 2*). The concentration of NE was not significantly different between the selected lines, but the ratio of EP/NE was significantly higher in the birds of the LGPS line (72.5% vs. 34.0%, $P < 0.01$, *Table 2*).

GENETIC SELECTION-INDUCED ALTERATIONS IN PLASMA CONCENTRATIONS OF CORTICOSTERONE

Genetic heritability related changes of blood concentration of CORT were measured using IRA, and the selected line differences were presented in the *Table 2*. The mean blood concentrations of CORT were 1.84 ± 0.19 and 1.49 ± 0.21 ng/mL for the KGB and LGPS birds, respectively (*Table 2*). There was a trend showing a higher blood concentration of CORT in the KGB birds than in the LGPS birds.

Discussion

The results of the present study provide evidence that selection for group productivity and longevity with reducing cannibalism and flightiness in birds results in altering regulation of the neurochemical system, including changes in blood concentrations of 5-HT, DA, EP, NE, and CORT. The data is consistent with previous findings in mammalian research that domestication of animals is associated with hereditary reorganization of the neuroendocrine system (Naumenko *et al.*, 1987, 1989; de Kloet *et al.*, 1996; Ferris, 2000) and changes of neurochemical homeostasis (Bilzard *et al.*, 1983; Crusio, 1996). The present findings further support our hypothesis that positive changes in birds' domestic behaviours and productive performances resulting from selective breeding likely reflect changes in the link between the nervous and endocrine systems. That is, when selection pressure is applied to productivity and survival as well as associated domestic behaviours by reducing cannibalism and flightiness, the pressure is simultaneously applied to the neuroendocrine system, resulting in changes of physiologic functions that control coping ability such as improving survivability and productivity.

SEROTONIN AND SELECTION FOR HIGH PRODUCTIVITY AND LONGEVITY WITH REDUCING CANNIBALISM

There were different levels of serotonin concentrations between the lines selected for high or low productivity and survivability. The results are consistent with the hypotheses that serotonin has multifunctional roles in modulating many types of behavioural and physiological processes, including feeding, sexual and aggressive behaviour (Mench and

Shea-Moore, 1995; Weiger, 1997; Olivier *et al.*, 1998). Abnormalities of blood and brain 5-HT and its metabolite, 5-HIAA, as well as the density of its receptors have been used as major indicators of alterations in behavioural adaptability and reproduction (Bell and Hobson 1994; Dinan, 1996; Popova *et al.*, 1997; Maswood *et al.*, 1998; Abumaris *et al.*, 2006; Nomura and Nomauro, 2006).

Serotonin functions as an inhibitory factor of aggression in controlling domestic behaviour (Popova *et al.*, 1975). Depletion or decrease of 5-HT concentration and ratio of 5-HIAA/5-HT in the central nervous system (CNS) have been implicated in dysfunctional behaviours, including aggressiveness and violence in human and non-human animals (Higley *et al.*, 1996; Unis *et al.*, 1997; Parmigiani *et al.*, 1999; Popova *et al.*, 1999; Gollan *et al.*, 2005; Popova, 2006; Wrase *et al.*, 2006), and cannibalism in birds (Barofsky *et al.*, 1983). However, biological roles of blood 5-HT in behavioural adaptation and motivational regulation are unclear. Decreased, increased and unchanged blood 5-HT concentrations have been found in association with behavioural dysfunctions, including aggressiveness (Cook *et al.*, 1995; Hanna *et al.*, 1995; Moffitt *et al.*, 1998). The conflicting data from different investigations could be related to different genetic selection programmes, species, behavioural evaluations and stressors used as well as duration and frequency of stressor presentation. Our results that higher blood 5-HT level associated with lower survivability resulting from higher cannibalism and flightiness in the LGPS birds is consistent with our hypothesis that different patterns of secreting of blood 5-HT are associated with specific behavioural parameters. Blood 5-HT level, positively correlated with aggressiveness, were also found in adolescents with behavioural conduct disorder (Unis *et al.*, 1997), borderline mental retardation patients with impaired impulse control and stress-induced aggression (Lenders *et al.*, 1998), and dominant male monkeys (Steklis *et al.*, 1986; Raleigh *et al.*, 1991). The association between 5-HT levels and aggressive behaviours supports our view that blood concentration of 5-HT could serve as a physiological indicator of the animal's coping ability to stress as well as a biological trait marker for domestic behaviours. In support of this view, other studies have determined that blood 5-HT levels are under genetic regulation (Jernej and Cicin-Sain, 1990; Cook and Leventhal, 1996), and its levels have been used as a stable marker in monitoring the course of certain dysfunctional behaviours (Stahl *et al.*, 1982), such as higher platelet 5-HT concentrations in schizophrenic patients (Muck-Seler *et al.*, 1999) and autism (Cook and Leventhal, 1996). Values of blood 5-HT also represent a heritable stable biological parameter in rodents (Jernej and Cicin-Sain 1990). As such rats, reverse selected for high and low platelet serotonin levels, were not affected by periodic oscillations, sex or age.

In the periphery, all 5-HT is stored in platelets (Anderson *et al.*, 1987; Jernej and Cicin-Sain, 1990), and platelet 5-HT is exchangeable with blood 5-HT but not brain 5-HT (Pietraszek *et al.*, 1992; Cook and Leventhal, 1996). Since the method used to measure 5-HT levels in platelets is to measure it in whole blood (Cook and Leventhal, 1996) and more than 99% of whole blood 5-HT is contained in platelets (Anderson *et al.*, 1987; Martin and Artigas, 1992), the two can be considered as belonging to a single compartment (Cook and Leventhal, 1996; Jernej *et al.*, 2000). Previous studies have shown that 5-HT systems between blood platelets and serotonergic neurons share some of anatomical and functional similarities (Stahl, 1977; Yan *et al.*, 1993), such as the active carrier mechanism, intracellular storage, receptors, and binding sites (Briley *et al.*, 1980; Paul *et al.*, 1980; Pletscher, 1988), platelets have been proposed as a peripheral tool for observation of central 5-HT system. However, the possibility of using platelets as a model system for brain neurons remains unclear. Previous studies have revealed that 5-HT levels in the CNS and peripheral tissues are genetically regulated differently (Popova, 1978; Lampagnani, 1986; Pietraszek *et al.*, 1992), and there are no correlations of either MAO activity (Young *et al.*, 1986; Virkkunen and Linnoila 1997) or 5-HT levels between brain

and platelet or blood (Popova, 1978; Takada *et al.*, 1996). Since 5-HT can not pass the brain-blood-barrier (Pietraszek *et al.*, 1992), blood and brain could be assumed as two compartments. About 90% of cerebral 5-HIAA effluxes directly into the blood (Pietraszek *et al.*, 1992) but there is no correlations between the changes in the CSF 5-HIAA and brain 5-HT (Martin and Artigas, 1992). These differences might be related to different mechanisms involve in regulating the uptake, synthesis and release of 5-HT between platelets and neurons (Chou *et al.*, 1983). For example, the binding potential (B_{max}/K_d) of 5-HT 2A receptors on platelets was not correlated with that of brain, and changes of 5-HT_{2A} receptors in platelets may not be a sign of similar changes in the CNS (Cho *et al.*, 1999). Ongoing studies in our laboratory aim to determine whether there are different cellular mechanisms underlying selection pressure-induced changes of 5-HT systems between periphery and brain in the presently selected lines.

The finding that LGP birds, compared to KGB birds, had higher concentrations of circulating 5-HT but lower productivity is consistent with the reports that 5-HT has a tonic, inhibitory effect on sexual behaviour and reproduction (Sirotkin and Schaeffer, 1997), such as inhibition of luteinising hormone (LH) secretion and ovulation (Nagatsuka, 1983; Morello *et al.*, 1992; Lorrain *et al.*, 1998). However, the roles of 5-HT in modulating sexual behaviour in female has been re-evaluated (Mendelson 1992), and is concluded that 5-HT can either inhibit or facilitate gonadotropins secretion, sexual hormone release, and sexual behaviour, which depends on which brain regions are involved (Nagatsuka, 1983; Gonzalez *et al.*, 1997; Wolf *et al.*, 1998), and what subtypes of 5-HT receptors are activated (Mendelson, 1992; Maswood *et al.*, 1998). As such effects of 5-HT on sexual behaviour were positively correlated to the stimulation of the pre-optic area (POA) and median eminence but negatively correlated to the stimulation of the ventromedial nucleus (VMN), zona incerta and arcuate nucleus (Gonzalez *et al.*, 1997). 5-HT₂ receptors in the POA were involved in the stimulatory function of 5-HT, while 5-HT_{1A} receptors mediated the inhibitory role in the VMN. Systemic injection of the selective 5-HT₂ and 5-HT₁ agonists, MK 212 and Ru 24969, sexual behaviour was facilitated and inhibited, respectively (Wilson and Hunter 1985). In addition, inhibited sexual behaviour was appeared by peripherally administrated 5-HT₂ antagonist, pirenperone (Mendelson and Gorzalka, 1985), and its effect was attenuated by the 5-HT agonist, quipazine. At present, cellular mechanisms that hereditarily regulate productivity between the selected lines are unclear. A further study has been set up in our lab to investigate selection pressure-induced neuronal morphofunctional plasticity, including 5-HT system, associated with different reproductive capability of KGB and LGPS birds.

DOPAMINE AND SELECTION FOR HIGH PRODUCTIVITY AND LONGEVITY WITH REDUCING CANNIBALISM

At physiological condition, DA is released centrally as a neurotransmitter, and is processed peripherally as a precursor of NE (Kuchel, 1991). Since DA cannot cross the blood-brain barrier (BBB), circulating DA could be either from adrenal cells or from leukocytes (Bergquist *et al.*, 1998). However, during pathological condition, DA can be released from adrenomedullary chromaffin cells, and the adrenal glands can be transformed from an adrenergic to a dopaminergic source (Snider and Kuchel, 1983). Abnormalities in the blood and brain DA systems have been associated with dysfunctional behaviours as well as with declined coping ability with stress (Driscoll *et al.*, 1998; Kuikka *et al.*, 1998; van Erp and Miczek, 2006; Watt *et al.*, 2006). In agreement with such findings, the selected lines under a genetically selected pressure conducted with a social stressor resulted in bi-directional changes of both blood DA concentrations and coping abilities. KGB birds, selected for higher productivity and longevity with reducing cannibalism and flightiness, but not LGPS birds, had significantly lower blood level of

DA, and better and faster adaptation to various stressors (Hester *et al.*, 1996a, b). Our findings suggest that heritability variations for DA concentrations reflect individual coping strategies (Benus *et al.*, 1991a, b; Vogel and Harris, 1991; Driscoll *et al.*, 1998). Similar to our findings, DA concentrations (Dimsdale and Ziegler 1991; Hjemdahl, 1993; Rolih and Ober, 1995), and phenotype (Nikulina and Kapralova, 1991; Vadasz *et al.*, 1992; de Kloet *et al.*, 1996) have been used as indicators to evaluate the capability of animals in coping with stressors and well being. In the bi-directionally selected Roman high (RHA/Verh) and low (RLA/Verh) avoidance rats, increased DA activity occurred in the RHA/Verh rats, associated with increased locomotor activity, a marker of anxiety and stressful status (Driscoll *et al.*, 1990; Corda *et al.*, 1997). Similar results have been reported in rats exposed to uncontrollable electric tail-shock (Scott *et al.*, 1996). In addition, rats selected for swim-test resistance showed higher concentrations of DA compared to rats selected for swim-test susceptible, these higher levels of DA were correlated with higher struggling activity and larger decrease in home-cage ambulatory activity and water intake after shock. Based on the present and previous observations, the selection-induced differences in dopaminergic and behavioural responsiveness are consistent with an interpretation that coping strategies of selected breeds are based on the inheritance pattern and phenotypic correlations of behavioural, physiological and neuroendocrine variables (Castanon *et al.*, 1995).

DA has also been thought to be involved in controlling domestic behaviours (Kuikka *et al.*, 1998; Korzan *et al.*, 2006). In CNS, increased DA concentration was found in selected brain regions in both humans and rodents following aggressive or defensive activities (Lewis *et al.*, 1994; Miczek *et al.*, 1994; Kuikka *et al.*, 1998; Mersmann, 1998; Volavka *et al.*, 2004; van Erp and Miczek, 2006). Higher levels of DA were also found in the brain of Japanese quails with aggressive behaviour (Edens, 1987). In the periphery, dysfunctions of DA were also involved in aggressive behaviours of patients with mental deficiency (Marazziti *et al.*, 1996). Similarly, higher baseline blood DA levels in our selected LGPS birds could be associated with the reorganization of behaviours, *i.e.*, higher cannibalism and flightiness, resulting in higher mortality. In contrast, the lower baseline of blood DA in our KGB birds associated with sedate and passive behaviours, as well as lower cannibalism and flightiness. Our findings suggest that genetic selection pressures directly and indirectly influences regulation of the brain dopaminergic system, and turn the activation of the DA system into a favour to survival behaviour in KGB birds. Genotypic dependence of animals in controlling domestic behaviour has also been demonstrated in other species. Sgoifo and co-investigators (1996) reported aggressive rats with higher activity of catecholamine system and short attack latency. Similarly, other studies investigated changes in the pharmacological profile of DA system with dysfunctional behaviours. Footshock induced aggression in mice can be facilitated by DA administered centrally, and L-dopa, a DA precursor, administered peripherally (Datla *et al.*, 1992). Benus and co-investigators (1991a) also reported that aggressive male mice were more sensitive to apomorphine, a D2 receptor agonist, than non-aggressive male. Biting behaviour (Shaikh *et al.*, 1991), footshock induced aggression (Datla *et al.*, 1992), and fighting (Ossowska *et al.*, 1996) can be significantly facilitated by apomorphine, but were suppressed by spiperone (Shaikh *et al.*, 1991; Datla *et al.*, 1992) and raclopride (Dennis *et al.*, 2006), D2 receptor antagonists, and by the dopamine blockers, haloperidol and pimozide (Nikulina and Kapralova, 1992; Pant and Nath, 1993; De Leon, 1994). Data from these studies suggest that DA plays a significant role in controlling aggressive behaviour.

Birds of LGPS line had a lower productivity but higher blood concentrations of DA compared to counterparts of KGB line. These results are consistent with the hypothesis that dopaminergic system is one of the main inhibitory neuronal systems that control the

development of the reproductive systems (Becu-Villalobos and Libertun, 1995) and productivity (Sotowska-Brochocka *et al.*, 1994). Although identification of mechanisms that underlie the inhibitory effects of DA on productivity in present lines is unclear, previous experimental findings suggest that the regulation might be related to genetic selection pressure induced changes in physiological functions of the neuroendocrine system, including the HAP and hypothalamic-pituitary-gonadal (HAG) axes. Provided studies had shown that endogenous DA secreted in the hypothalamus exhibited catecholamine's tonic inhibition of Luteinising hormone-releasing hormone (LHRH) release (Contijoch *et al.*, 1992) and suppression LH secretion (Martin *et al.*, 1981). In rodents, the biological functions of DA on productivity had been confirmed by administering exogenous DA that significantly reduced the LHRH level from the hypothalamus and lowered LH secretion from the pituitary (Tasaka *et al.*, 1985). Similar results were obtained from poultry studies. In hens, increase in dopaminergic activity inhibited the release of LH by exerting an inhibitory influence on the LHRH containing cells in the hypothalamus (Macnamee and Sharp 1989). The effects of DA could be related to increase in tyrosine hydroxylase activity (Thiery *et al.*, 1995) and activity of DA receptors (Andersson *et al.*, 1988). The later had been verified by either administering of DA receptor agonist 2-Br-alpha-ergocryptine to lower plasma LH levels (Owens *et al.*, 1980) or blocking DA inhibitory effect by use of DA receptor blocker halperidol (Tasaka *et al.*, 1985).

Although it is unclear what mechanisms underlie the regulation of release DA peripherally in the presently selected birds, previous studies have reported that such release could be induced by stress-related unbalance of activities of enzymes involved in catecholamine metabolism (Kuchel, 1991). Stressor stimulates TH activity and, at the same time, inhibits activities of the enzyme dopamine- β -hydroxylase (D β H) that converts DA to NE (Kuchel *et al.*, 1982). If D β H activation cannot keep pace with activation of TH, DA synthesis predominates, and extra DA is released peripherally. Without detecting TH activities in the presently selected lines, we can not assume that LGPS birds have a higher TH activation. However, previous studies have shown that TH activation can be increased by selection for domestic behaviour in animals (Dygalo *et al.*, 1988; Kulikov *et al.*, 1989), and that hereditary based stress triggers different regulations of gene expressions for catecholamine biosynthetic enzymes in the adrenal medulla (Nankova and Sabban, 1999).

EPINEPHRINE AND NOEPINEPHRINE AND SELECTION FOR HIGH PRODUCTIVITY AND LONGEVITY WITH REDUCING CANNIBALISM

EP is released almost exclusively from the adrenal medulla, with a small amount synthesized in the brain, while NP is released from sympathetic neurons, with significant amount synthesized in the CNS, especially, in the hypothalamus (Bullock *et al.*, 1995). As "stress hormones", both EP and NE participate in a number of physiologic and pathologic processes, including regulation of emotion and motivation in response to stimulations. Changes in EP and NE levels, as well as the ratio of EP/NE have been used as indicators of the 'organisms' well-being and capability of cope to stress (Goldstein, 1981). In response to stimulations from selected pressure, the lower blood concentration of EP and lower ratio of circulating EP/NE found in KGB birds is consistent with the multifunctional roles of EP and NE. In contrast, the LGP birds had a higher blood concentration of EP and higher ratio of EP/NE, which correlates with their lower productivity and higher mortality from cannibalism. Furthermore, the ratio of EP/NE has been used as an indicator of well-being. A higher EP/NE ratio had been found in severely ill patients (Lechin *et al.*, 1996), stressful states (Vollmer, 1996; Piercecchi-Marti *et al.*, 1999), and dominant male monkeys (Dillon *et al.*, 1992). Collectively, these data support our hypothesis that genotypic or phenotypic variation of the EP/NE, in response to selection, may reflect

individual coping strategies that control changes in the animal's behavioural patterns.

In the present study the major factor causing the EP/EN increase in LGP birds was higher concentration of EP, since there was not significantly different in the concentrations of NE between the lines. A similar upregulation of EP concentration by increased biosynthesis was found in turkeys selected for higher adrenal response to cold stress (HL) (Brown and Nestor, 1974). Birds of the HL line also, similar to the LGP birds, laid significantly less eggs and had hyperactivity and poorer feeding efficiency. The present and previous results support the notion that genetic selection for various physiological indexes could have altered the effects of the central and peripheral dopaminergic system in an attempt to adapt to stress. It could mean that EP/NE can be used as an indicator for selecting and evaluate the animal's well-being.

CORTICOSTERONE AND SELECTION FOR HIGH PRODUCTIVITY AND LONGEVITY WITH REDUCING CANNIBALISM

Corticosterone is released from the adrenal cortex, the rate of its secretion being controlled by activation of the hypothalamic-pituitary-adrenal (HPA) axis via a double negative feedback loop. During pathologic conditions, such as under various stressors including social stress, concentration of CORT can be increased by overriding the HPA control system. As one of the "stress hormones", CORT has multifunctional roles in both normal and abnormal states, including regulation of the organism's behavioural patterns, coping styles and well-being.

Although differences in the baseline concentration of CORT were not found here to be significant, there was a tendency for CORT concentration to be higher in KGB birds than in those of the reverse selected line. The heritably different levels of CORT between the present selected lines could be the results of genetically different regulations in CORT secretion as induced by selected pressure. Supporting our findings, the value of circulating CORT has been used as a genetic selection criterion for evaluation of coping ability in responsiveness to stress, such as in open-field exposure (Jones *et al.*, 1992), physical and manual restraint (Mills and Faure, 1991; Jones and Satterlee, 1996; Savory and Mann, 1997), fear of humans (Jones *et al.*, 1994), cold and heart temperature stimulation (Brown and Nestor, 1974; Hester *et al.*, 1996a), and resistance to infection and inflammation (Gross and Colmano, 1971). Based on these observations, changes of CORT levels associated with different behaviours are a hereditary hallmark of the organism's responsiveness to stressors, and effects of CORT on stress-induced organism's responsiveness may depend on its levels, *i.e.*, a level slightly or excessively changed from its physiological basis, which could protect or destruct the organism's coping capability (Sapolsky and Meaney, 1986; Jones and Satterlee, 1996). The higher CORT concentration in KGB birds than in LGPS birds (1.84 vs. 1.49 ng/ml, $P=0.1$) might be at the high end of physiologic levels. Higher baseline levels of CORT signal a slight hyperactivity of the CORT system in KGB birds compared that in LGP birds, which is consistent with the previous findings that KGB birds had greater adrenal weights compared to those of control line (Hester *et al.*, 1996c). This view is further supported by observations in domestic Larger White pigs (Fernandez *et al.*, 1994; Blanchard *et al.*, 1993) and aggressive rodents (Korte *et al.*, 1996). As such a tendency to higher baseline plasma cortisol levels was found in subordinated pigs and in mice with longer attack latencies as compared with those aggressive counterparts, respectively. In addition, Shively and Kaplan (1984) determined that the subordinate Macaca fascicularis had heavier adrenal glands with higher plasma glucose levels but less atherosclerosis than those of dominants. The above findings suggest that upregulation of the adrenal system in KGB birds is part of a defence mechanism against environmental challenges, which may underlie their higher survivability, sedate and passive behaviours, and better coping to social, handling and

environmental stressors (Hester *et al.*, 1996a, b; Muir and Craig, 1996). Without stress challenges, we can not assume that KGB birds have a lower reactivation of CORT system in responses to stimuli. However, Sapolsk (1990) reported that, compared to high ranking adult male baboons, low ranking males had a higher basal cortisol and lower cortisol in response to the challenges of capture and corticotropin-releasing factor (CRF). Collectively, these results are consistent with the notion that adrenal hypertrophy may be an indicator of greater adaptation to stress (Siegel, 1971). Therefore, alterations in regulation of adrenocortical activity appear to be essential in controlling of coping activity and styles, as well as mobilization of body energy sources such as glycogen and triglyceride in response to stressors (Gaillard and Al-Damluji, 1987; Tempel and Leibowitz, 1994).

CORT defensive response to stress in KGB birds could be associated also with its role as an immunomodulator. As such following challenges with *Escherichia coli*, infection can be prevented by pre-treatment with CORT (Gross and Colmano, 1970). CORT can also protect the organism from damage by an excessive response of its immune system (Gross and Colmano, 1970; Munck *et al.*, 1984). Our data and others are consistent with the notion that, under physiologic condition, one of the primary roles of endogenous glucocorticoids may be immunomodulation rather than immunosuppression (Dhabhar *et al.*, 1994); and that small increases in levels of glucocorticoid may enhance immune functions (Stanulis *et al.*, 1997). Thus in one of our parallel studies has found that KGB birds had an inherited basis of higher cellular-modulated immunity than that of LGPS birds (Chen *et al.*, 2000). In contrast, LGPS birds had higher levels of circulating immunoglobulin G and higher ratios of circulating Heterophil/lymphocyte. The different inherited characteristics of immunity could be associated with better and faster adaptability to various stressors as reported previously (Hester *et al.*, 1996a, b). In addition, except for the direct effects of CORT on animal behaviours such as aggression (Bonson *et al.*, 1994; Kruk *et al.*, 1998), the sedate and passive behaviours of the KGB birds could also related to indirectly effects through the regulating serotonergic system (Chaouloff, 1995; Joels *et al.*, 1995). Thus, the effects of CORT may enhance brain serotonergic turnover (Summers *et al.*, 2000), increasing 5-HT formation from tryptophan (Millard *et al.*, 1972), and affect the activity of 5-HT receptors (Mendelson and McEwen, 1992). Collectively, these results are consistent with the protective functions of CORT and could be, at least partly, related to natural mechanisms permitting animals to remain alert and keep physiological homeostasis in response to relevant stimulants.

Conversely, CORT could have inhibitory effects on the animal's coping ability, and negative influence physical parameters have been reported. For example, there were approximately 2-3 folds significantly higher concentrations of CORT in birds genetically selected for high adrenal response (HL) in response to mechanical restraint, compared to that in birds of the low response line (Brown and Nestor, 1974). Birds of the HL failed to maintain physiologic homeostasis, including slower growth (Brown *et al.*, 1959), overstated fear (Jones *et al.*, 1992; Jones and Satterlee, 1996), were higher excitable (Brown and Nestor, 1973, 1974), laid fewer eggs, and had higher natural mortality (Brown and Nestor, 1974).

A higher productivity and a tendency for CORT concentration to be higher in KGB birds compared to those of LGPS line suggest that CORT may have functions in regulating productivity of hens. The hypothesis is consistent with previous findings that the ovary is the one of the most probable target tissue for CORT (Etches *et al.*, 1983). CORT may exert its ovulation-inducing effect directly on the mature follicle (Etches *et al.*, 1984; Lang *et al.*, 1984) through an anatomical juxtaposition of the left ovary and adrenal gland (Etches *et al.*, 1984), or may alter the responsiveness of ovarian tissue to gonadotropins through modulating the reaction of the hypothalamus to tropic stimuli (Etches *et al.*, 1984). A previous study has demonstrated that CORT involves in the preovulatory surge of LH

which results in the first egg of a sequence (Sharp and Beuving, 1978). On the other hand, effects of CORT on productivity may through affects the serotonergic, dopaminergic systems or both which have been known to be involved in regulation of reproduction (Summers *et al.*, 2000; also see above discussion). In addition, CORT has impact on inflammatory and immunological processes (Gaillard and Al-Damluji, 1987). At physiological levels CORT enhance immune functions (Stanulis *et al.*, 1997), and protects organisms from reaching excessive immune reactive levels to cause damage to itself (Munck *et al.*, 1984), or inhibit autoimmunity which has been identified as one of the major reasons of impaired productivity (Bacon, 1987). Furthermore, previous studies have evidenced that CORT has important effects on caloric intake. Under normal condition, endogenous CORT involves in controlling ingestion and metabolism (Tempel and Leibowitz, 1994), such as catabolic effects on carbohydrate, fat and protein to maintain glucose homeostasis. Whether higher blood concentration of CORT in KGB birds affect feed efficiency is unclear, previous study had shown that KGB birds fed more, at least at first three weeks during the initial adaptation to layer house cages, than those of commercial lines (Craig and Muir, 1996b). The better feeding behaviours of KGB birds may be associated with their better adaptability to a variety of stressors (Hester *et al.*, 1996a, b), and their higher productivity. Currently, it is unclear whether LPGS birds have more active adrenal system in synthesizing CORT in response to stress, but previous study in the Syracuse high- (SHA) and low-avoidance (SLA) strains of rats has shown that, in response to stress, the SHA rats with smaller adrenals had larger responses of the HPA axis (Brush *et al.*, 1991).

Taken together, these data suggest that the effect of CORT on the animal's coping ability is biphasic and dependent on its concentration, which results in its attempt to engage or avoid environmental challenges, *i.e.*, at physiological levels it can protect the organism to cope with stressors, whereas at pathological levels it results in declining coping ability. Stress-induced reactions of the CORT system may be dependent on the species, breeds and stressors used as well as duration and frequency of stressor presentation. Although conflicting results have been reported, there is consensus as to the cellular mechanisms which regulate CORT secretion, *i.e.*, mediators are from the HPA axis. Alterations in the levels of corticotrophin-releasing hormone (CRH) and adrenocorticotropin (ACTH) control CORT secretion from adrenal glands (Rots *et al.*, 1995). Our present data support the view that individual differences in coping styles is coherent with a set of functions of the neuroendocrine system that underlies behavioural characteristics (Sgoifo *et al.*, 1996).

In conclusion, the present study demonstrates that selection for high and low group productivity and longevity with alterations in cannibalism and flightiness affected the regulations of the neuroendocrine system of selected birds, and that CORT and monoamines, such as 5-HT, DA, EP and NE, were differently regulated by selection pressure. Compared to the reverse selected birds, KGB birds selected for higher productivity and longevity with higher domestic behaviours, *i.e.*, low cannibalism and flightiness, may have a positive alterations in the neuroendocrine system, *i.e.*, lower blood concentrations of 5-HT, DA and EP, and higher levels of plasma CORT, which may be associated with their better coping to a novel environment and greater resistance to stressors. The unique homeostatic characteristics of each selected line may provide a neurobiological basis for investigating effects of genetic factors on physiological functions of biogenic amines involved in productivity and longevity related to domestic behaviours. The present data and data from others suggest that blood 5-HT, DA and EP as well as ratio of EP/NE, can be used as physiological indicators for genetically selecting birds with great resistance to social stress to eliminating beak trimming. This finding could be extended to other farm animals for improving their well-being.

Acknowledgements

We thank J. Johnson for helping to collect samples. This work was supported by a USDA-NRI grant.

References

- APPLEBY, M.C. and HOGARTH, G.S. (1991) Welfare of laying hens in cages and alternative systems: environmental, physical and behavioural aspects. *World's Poultry Science Journal* **47**: 109-128.
- ABUMARIA, N., RYGULA, R., HAVEMANN-REINECKE, U., RUTHER, E., BODEMER, W., ROOS, C. and FLUGGE, G. (2006) Identification of genes regulated by chronic social stress in the rat dorsal raphe nucleus. *Cellular and Molecular Neurobiology* **26**: 145-162.
- ANDERSON, G.M., FEIBEL, F.C. and COHEN, D.J. (1987) Determination of serotonin in whole blood, platelet-rich plasma, platelet-poor plasma and plasma ultrafiltrate. *Life Science* **40**: 1063-1070.
- ANDERSSON, K., FUXE, K., ENEROTH, P., HARFSTRAND, A. and AGNATI, L.F. (1988) Involvement of D1 dopamine receptors in the nicotine-induced neuro-endocrine effects and depletion of diencephalic catecholamine stores in the male rat. *Neuroendocrinology* **48**: 188-200.
- BARRACLOUGH, C.A. (1992) Neural control of the synthesis and release of luteinizing hormone-releasing hormone. *Ciba Foundation symposium* **168**: 233-246.
- BACON, L.D. (1987) Influence of the major histocompatibility complex on disease resistance and productivity. *Poultry Science* **66**: 802-811.
- BALABAN, E., ALPER, J.S. and KASAMON, Y.L. (1996) Mean genes and the biology of aggression: a critical review of recent animal and human research. *Journal of Neurogenetics* **11**: 1-43.
- BARAKK, Y. and NASHIACH, M. (2004) Aggression – from basic science to real life. *Israel Medical Association Journal* **5**: 667-668.
- BARRACLOUGH, C.A. (1992) Neural control of the synthesis and release of luteinizing hormone-releasing hormone. *Ciba Foundation Symposia* **168**: 233-246.
- BAROFSKY, A.L., TAYLOR, J., TIZABI, Y., KUMAR, R. and JONES-QUARTEY, K. (1983) Specific neurotoxin lesions of median raphe serotonergic neurons disrupt maternal behaviour in the lactating rat. *Endocrinology* **113**: 1884-1893.
- BAYYARI, G.R., HUFF, W.E., RATH, N.C., BALOG, J.M., NEWBERRY, L.A., VILLINES, J.D., SKEELES, J.K., ANTHONY, N.B. and NESTOR, K.E. (1997) Effect of the genetic selection of turkeys for increased body weight and egg production on immune and physiological responses. *Poultry Science* **76**: 289-296.
- BECU-VILLALOBOS, D. and LIBERTUN, C. (1995) Development of gonadotropin-releasing hormone (GnRH) neuron regulation in the female rat. *Cellular and Molecular Neurobiology* **15**: 165-176.
- BELL, R. and HEPPER P.G. (1987) Catecholamines and aggression in animals. *Behavioural Brain Research* **23**: 1-21.
- BELL, R. and HOBSON, H. (1994) 5-HT_{1A} receptor influences on rodent social and agonistic behaviour: a review and empirical study. *Neuroscience and Biobehavioural Reviews* **18**: 325-338.
- BENUS, R.F., BOHUS, B., KOOLHAAS, J.M. and VAN OORTMERSSEN, G.A. (1991a) Behavioural differences between artificially selected aggressive and non-aggressive mice: response to apomorphine. *Behavioural Brain Research* **43**: 203-208.
- BENUS, R.F., BOHUS, B., KOOLHAAS, J.M. and VAN OORTMERSSEN, G.A. (1991b) Heritable variation for aggression as a reflection of individual coping strategies. *Experientia* **47**: 1008-1019.
- BERGQUIST, J., TARKOWSKI, A., EWING, A. and EKMAN, R. (1998) Catecholaminergic suppression of immunocompetent cells. *Immunology Today* **19**: 562-567.
- BERMAN, M.E. and COCCARO, E.F. (1998) Neurobiologic correlates of violence: relevance to criminal responsibility. *Behavioural Sciences and the Law* **16**: 303-318.
- BILZARD, D.A., FREEDMAN, L.S. and LIANG, B. (1983) Genetic variation, chronic stress, and the central and peripheral noradrenergic system. *American Journal of Physiology* **R 600**: 600-605.
- BLANCHARD, D.C., SAKAI, R.R., MCEWEN, B., WEISS, S.M. and BLANCHARD, J.R. (1993) Subordination stress: behavioural, brain, and neuroendocrine correlates. *Behavioural Brain Research* **58**: 113-121.
- BLUM, K. and SHERIDAN, P.J. (1995) Dopamine D2 receptor gene variants-association and linkage studies in impulsive-addictive-compulsive behaviour. *Pharmacogenetics* **5**: 121-141.
- BLUM, K., WOOD, R.C., BRAVERMAN, E.R., CHEN, T.J. and SHERIDAN, P.J. (1995) The D2 dopamine receptor gene as a predictor of compulsive disease: Bayes' theorem. *Functional Neurology* **10**: 37-44.
- BONSON, K.R., JOHNSON, R.G., FIORELLA, D., RABIN, R.A. and WINTER, J.C. (1994) Serotonergic control of androgen-induced dominance. *Pharmacology Biochemistry Behaviour* **49**: 313-322.

- BRILEY, M.S., LANGER, S.Z., RAISMAN, R., SECHTER, D. and ZARIFIAN, E. (1980) Tritiated imipramine binding sites are decreased in platelets of untreated depressed patients. *Science* **209**: 303-305.
- BROWN, K.I. (1959) 'Stress' and its implication in poultry production. *World's Poultry Science Journal* **15**: 255-263.
- BROWN, K.I. and NESTOR, K.E. (1973) Some physiological responses of turkeys selected for high and low adrenal response to cold stress. *Poultry Science* **52**: 1948-54
- BROWN, K.I. and NESTOR, K.E. (1974) Implications of selection for high and low adrenal response to stress. *Poultry Science* **53**: 1297-1306.
- BRUSH, F.R., ISAACSON, M.D., PELLEGRINO, L.J., RYKASZEWSKI, I.M. and SHAIN, C.N. (1991) Characteristics of the pituitary-adrenal system in the Syracuse high- and low-avoidance strains of rats (*Rattus norvegicus*). *Behavioural Genetics* **21**: 35-48.
- BULLOCK, J., BOYLE, J. III and WANG, M.B. (1995) Physiology. Williams & Wilkins. Malvern, PA.
- CASTANON, N., PEREZ-DIAZ, F. and MORMEDE, P. (1995) Genetic analysis of the relationships between behavioural and neuroendocrine traits in Roman high and low avoidance rat lines. *Behavioural Genetics* **25**: 371-384.
- CHAOULOFF, F. (1995) Regulation of 5-HT receptors by corticosteroids: where do we stand? *Fundamental and Clinical Pharmacology* **9**: 219-233.
- CHENG, H.W., DILLWORTH, G., SINGLETON, P., CHEN, Y. and MUIR, W.M. (2001a) Effect of genetic selection for productivity and longevity on blood concentrations of serotonin, catecholamine and corticosterone of chickens. *Poultry Science* **80**: 1278-1285.
- CHENG, H.W., EICHER, S.D., CHEN, Y., SINGLETON, P. and MUIR, W.M. (2001b) Effect of genetic selection for group productivity and longevity on immunological and hematological parameters of chickens. *Poultry Science* **80**: 1079-1086.
- CHENG, H.W. and MUIR, W.M. (2005) The effects of genetic selection for survivability and productivity on chicken physiological homeostasis. *World's Poultry Science Journal* **61**: 383-397.
- CHO, R., KAPUR, S., DU, L. and HRDINA, P. (1999) Relationship between central and peripheral serotonin 5-HT_{2A} receptors: a positron emission tomography study in healthy individuals. *Neuroscience Letter* **261**: 139-142.
- CHOU, D.T., CUZZONE, H. and HIRSH, K.R. (1983) Assessment of blood platelets as a model for CNS response: comparative effects of caffeine on 5-HT uptake and release mechanisms in rat platelets and rat brain serotonin neurons. *Journal of life science* **33**: 1149-56.
- COOK, E.H., STEIN, M.A. ELLISON, T., UNIS, A.S. and LEVENTHAL, B.L. (1995) Attention deficit hyperactivity disorder and whole-blood serotonin levels: effects of comorbidity. *Psychiatry Research* **57**: 13-20.
- COOK, E.H. and LEVENTHAL, B.L. (1996) The serotonin system in autism. *Current Opinion Pediatrics* **8**: 348-354.
- CONTIJOCH, A.M., GONZALEZ, C., SINGH, H.N., MALAMED, S., TRONCOSO, S. and ADVIS, J.P. (1992) Dopaminergic regulation of luteinizing hormone-releasing hormone release at the median eminence level: immunocytochemical and physiological evidence in hens. *Neuroendocrinology* **55**: 290-300.
- CORDA, M.G., LECCA, D., PIRAS, G., DI CHIARA, G. and GIORGI, O. (1997) Biochemical parameters of dopaminergic and GABAergic neurotransmission in the CNS of Roman high-avoidance and Roman low-avoidance rats. *Behavioural Genetics* **27**: 527-536.
- CRAIG, J.V. and SWANSON, J.C. (1994) Review: welfare perspectives on hens kept for egg production. *Poultry Science* **73**: 921-938.
- CRAIG, J.V. and MUIR, W.M. (1996a) Group selection for adaptation to multiple-hen cages: beak-related mortality, feathering, and body weight responses. *Poultry Science* **75**: 294-302.
- CRAIG, J.V. and MUIR, W.M. (1996b) Group selection for adaptation to multiple-hen cages: behavioural responses. *Poultry Science* **75**: 1145-1155.
- CRAIG, J.V., DEAN, W.F., HAVENSTEIN, G.B., KRUGER, K.K., NESTOR, K.E., PURCHASE, G.H., SIEGEL, P.B. and VAN WICKLEN, G.L. (1999) Guidelines for poultry husbandry. In: Guide for the care and use of agricultural animals in agricultural research and teaching. *Federation of Animal Science Societies* Pp.55-66.
- CRUSIO, W.E. (1996) The neurobehavioural genetics of aggression. *Behavioural Genetics* **26**: 459-461.
- DATLA, K.P., SEN, A.P. and BHATTACHARYA, S.K. (1992) Dopaminergic modulation of footshock induced aggression in paired rats. *Indian Journal Experimental Biology* **30**: 587-591.
- DAVIDSON, R.J., PUTNAM, K.M. and LARSON, C.L. (2000) Dysfunction in the neural circuitry of emotion regulation—a possible prelude to violence. *Science* **289**: 591-594.
- DE KLOET, E.R., KORTE, S.M., ROTS, N.Y. and KRUK, M.R. (1996) Stress hormones, genotype, and brain organization. Implications for aggression. *Annals of the New York Academy Sciences* **794**: 179-191.
- DE LEON, O.A. (1994) The neurobiological bases of aggression: the pharmacotherapeutic implications. *Revista médica de Panama* **19**: 106-116.
- DENNIS, R., ZHANG, H.M., BACON, L.D., ESTEVEZ, I. and CHENG, H.W. (2004) Behavioural and physiological features of chickens diversely selected for resistance to Avian Disease. 1. Selected inbred lines differ in behavioural and physical responses to social stress. *Poultry Science* **83**: 1489-96.

- DENNIS, R., MUIR, W.M. and CHENG, H.W. (2006) Effects of raclopride on aggression and stress in diversely selected chicken lines. *Behavioural Brain Research* **175**: 104-111.
- DHABHAR, F.S., MILLER, A.H., STEIN, M., MCEWEN, B.S. and SPENCER, R.L. (1994) Diurnal and acute stress-induced changes in distribution of peripheral blood leukocyte subpopulations. *Brain Behaviour and Immunity* **8**: 66-79.
- DIETL, M.M. and PALACIOS, J.M. (1988) Neurotransmitter receptors in the avian brain. I. Dopamine receptors. *Brain Research* **439**: 354-359.
- DILLON, J.E., RALEIGH, M.J., MCGUIRE, M.T., BERGIN-POLLACK, D. and YUWILER, A. (1992) Plasma catecholamines and social behaviour in male vervet monkeys (*Cercopithecus aethiops sabaeus*). *Physiology and Behaviour* **51**: 973-977.
- DIMSDALE, J.E. and ZIEGLER, M.G. (1991) What do plasma and urinary measures of catecholamines tell us about human response to stressors? *Circulation* **83**: 1136-42.
- DINAN, T.G. (1996) Serotonin: current understanding and the way forward. *International Clinical Psychopharmacology Suppl* **1**: 19-21.
- DOHMS, J.E. and METZ, A. (1991) Stress—mechanisms of immunosuppression. *Veterinary Immunology and Immunopathology* **30**: 89-109.
- DRISCOLL, P., DEDEK, J., D'ANGIO, M., CLAUSTRE, Y. and SCATTON, B. (1990) A genetically-based model for divergent stress responses, behavioural, neurochemical and hormonal aspects. In: Pliska, V. and G. Stranzinger, eds. *Farm animals in biomedical research*. Hamburg: Verlag Paul Parey. Pp.97-107.
- DRISCOLL, P., ESCORIHUELA, R.M., FERNANDEZ-TERUEL, A., GIORGI, O., SCHWEGLER, H., STEIMER, T., WIERSMA, A., CORDA, M.G., FLINT, J., KOOLHAAS, J.M., LANGHANS, W., SCHULZ, P.E., SIEGEL, J. and TOBENA, A. (1998) Genetic selection and differential stress responses. The Roman lines/strains of rats. *Annals of the New York Academy Sciences* **851**: 501-510.
- DUNCAN, I.J. (1998) Behaviour and behavioural needs. *Poultry Science* **77**:1766-1772.
- DYGALO, N.N., BYKOVA, T.S. and NAUMENKO, E.V. (1988) Brain tyrosine hydroxylase activity in behaviour-selected silver foxes. *Zhurnal evoliutsionno< biokhimii i fiziologii* **24**: 503-508.
- EDEND, F.W. (1987) Agonistic behaviour and neurochemistry in grouped Japanese quail. *Comparative Biochemistry and Physiology A* **86**: 473-479.
- ELDAGHAYES, I., ROTHWELL, L., WILLIAMS, A., WITHERS, D., BALU, S., DAVISON, F. and KAISER, P. (2006) Infectious bursal disease virus: strains that differ in virulence differentially modulate the innate immune response to infection in the chicken bursa. *Viral Immunology* **19**: 83-91.
- ETCHES, R.J., MACGREGOR, H.E., MORRIS, T.F. and WILLIAMS, J.B. (1983) Follicular growth and maturation in the domestic hen (*Gallus domesticus*). *Journal of Reproduction Fertility* **67**: 351-358.
- ETCHES, R.J., WILLIAMS, J.B. and RZASA, J. (1984) Effects of corticosterone and dietary changes in the hen on ovarian function, plasma LH and steroids and the response to exogenous LH-RH. *Journal of Reproduction Fertility* **70**: 121-130.
- FERNANDEZ, X., MEUNIER-SALAUN, M.C. and MORMEDE, P. (1994) Agonistic behaviour, plasma stress hormones, and metabolites in response to dyadic encounters in domestic pigs: interrelationships and effect of dominance status. *Physiology and Behaviour* **56**: 841-847.
- FERRIS, C.F. (2000). Adolescent stress and neural plasticity in hamsters: a vasopressin-serotonin model of inappropriate aggressive behaviour. *Experimental Physiology* **85**: 85S-90S.
- FRASER, D. and RUSHEN, J. (1987) Aggressive behaviour. *Veterinary Clinics of North America-Food Animal Practice* **3**: 285-305.
- GAILLARD, R.C. and AL-DAMLUJI, S. (1987) Stress and the pituitary-adrenal axis. *Baillieres Clinical Endocrinology Metabolism* **1**: 319-354.
- GEORGES, M. (1999) Towards marker assisted selection in livestock. *Reproduction Nutrition Development* **39**: 555-561.
- GLATZ, P.C. (2005) Beak trimming handbook. A report for the Australian poultry cooperative research centre. Poultry CRC Publication No 03-22.
- GOLAN, J.K., LEE, R. and COCCARO, E.F. (2005) Developmental psychopathology and neurobiology of aggression. *Development and Psychopathology* **17**: 1151-1171.
- GOLDSTEIN, S.D. (1981) Plasma norepinephrine as an indicator of sympathetic neural activity in clinical cardiology. *American Journal Cardiology* **48**: 1147-1154.
- GONZALEZ, M.I., GREENGRASS, P., RUSSELL, M. and WILSON, C.A. (1997) Comparison of serotonin receptor numbers and activity in specific hypothalamic areas of sexually active and inactive female rats. *Neuroendocrinology* **66**: 384-392.
- GOODSON, J.L. (2005) The vertebrate social behaviour network: evolutionary themes and variations. *Hormones and Behaviour* **48**: 11-22.
- GROSS, W.B. and COLMANO, G. (1970) Corticosterone and ACTH as treatments for *Escherichia coli* infection in chickens. *Poultry Science* **49**: 1256-1258.
- GROSS, W.B. and COLMANO, G. (1971) Effect of infectious agents on chickens selected for plasma corticosterone response to social stress. *Poultry Science* **50**: 1213-1220.

- HALLER, J., MAKARA, G.B. and KRUK, M.R. (1997) Catecholaminergic involvement in the control of aggression: hormones, the peripheral sympathetic, and central noradrenergic systems. *Neuroscience and Biobehavioural reviews* **22**: 85-97.
- HANNA, G.L., YUWILER, A. and COATES, J.K. (1995) Whole blood serotonin and disruptive behaviours in juvenile obsessive-compulsive disorder. *Journal of the American Academy of Child and Adolescent Psychiatry* **34**: 28-35.
- HESTER, P.Y., MUIR, W.M., CRAIG, J.V. and ALBRIGHT, J.L. (1996a) Group selection for adaptation to multiple-hen cages: production traits during heat and cold exposures. *Poultry Science* **75**: 1308-1314.
- HESTER, P.Y., MUIR, W.M. and CRAIG, J.V. (1996b) Group selection for adaptation to multiple-hen cages: humoral immune response. *Poultry Science* **75**: 1315-1320.
- HESTER, P.Y., MUIR, W.M., CRAIG, J.V. and ALBRIGHT, J.L. (1996c) Group selection for adaptation to multiple-hen cages: hematology and adrenal function. *Poultry Science* **75**: 1295-1307.
- HIGLEY, J.D., KING, S.T., HASERT, M.F., CHAMPOUX, M., SUOMI, S.J. and LINNOILA, M. (1996) Stability of interindividual differences in serotonin function and its relationship to severe aggression and competent social behaviour in Rhesus macaque females. *Neuropsychopharmacology* **14**: 67-76.
- HJEMDAHL, P. (1993) Plasma catecholamines—analytical challenges and physiological limitations. *Baillieres Clinical Endocrinology Metabolism* **7**: 307-53.
- JERNEJ, B. and CICIN-SAIN, L. (1990) Platelet serotonin level in rats is under genetic control. *Psychiatry Research* **32**: 167-174.
- JERNEJ, B., BANOVIC, M., CICIN-SAIN, L., HRANILOVIC, D., BALIJA, M., ORESKOVIC, D. and FOLNEGOVIC-SMALC, V. (2000) Physiological characteristics of platelet/circulatory serotonin: study on a large human population. *Psychiatry Research* **94**: 153-162.
- JOELS, M., HESEN, W. and de KLOET, E.R. (1995) Long-term control of neuronal excitability by corticosteroid hormones. *The Journal of steroid biochemistry and molecular biology* **53**: 315-323.
- JONES, R.B., SATTERLEE, D.G. and RYDER, F.H. (1992) Research note: open-field behaviour of Japanese quail chicks genetically selected for low or high plasma corticosterone response to immobilization stress. *Poultry Science* **71**: 1403-1407.
- JONES, R.B., MILLS, A.D., FAURA, J.M. and WILLIAMS, J.B. (1994) Restraint, fear, and distress in Japanese quail genetically selected for long or short tonic immobility reactions. *Physiology and Behaviour* **56**: 529-534.
- JONES, R.B. and SATTERLEE, D.G. (1996) Threat-induced behavioural inhibition in Japanese quail genetically selected for contrasting adrenocortical response to mechanical restraint. *British poultry science* **37**: 465-70.
- KANNAN, G., HEATH, J.L., WABECK, C.J., SOUZA, M.C., HOWE, J.C. and MENCH, J.A. (1997) Effects of crating and transport on stress and meat quality characteristics in broilers. *Poultry Science* **76**: 523-529.
- KAPPE, S.M. (1999) Utilization of gene mapping information in livestock animals. *Theriogenology* **51**: 135-147.
- KILGOUR, R.J. and SZANTAR-CODDINGTON, M.R. (1997) The arena test and cortisol response of sheep as indirect selection criteria for the improvement of lamb survival. *Animal reproduction science* **46**: 97-108.
- KOOLHAAS, J.M., KORTE, S.M., DE BOER, S.F., VAN DER VEGT, B.J., VAN REENEN, C.G., HOPSTER, H., DE JONG, I.C., RUIS, M.A. and BLOKHUIS, H.J. (1999) Coping styles in animals: current status in behaviour and stress-physiology. *Neuroscience and biobehavioural reviews* **23**: 925-935.
- KOPIN, I.J. (1984) Avenues of investigation for the role of catecholamines in anxiety. *Psychopathology Suppl* **1**: 83-97.
- KORTE, S.M., MEIJER, O.C., DE KLOET, E.R., BUWALDA, B., KEIJSER, J., SLUYTER, F., VAN OORTMERSEN, G. and BOHUS, B. (1996) Enhanced 5-HT_{1A} receptor expression in forebrain regions of aggressive house mice. *Brain Research* **736**: 338-343.
- KORZAN, W.J., FORSTER, G.L., WATT, M.J. and SUMMERS, C.H. (2006) Dopaminergic activity modulation via aggression, status, and a visual social signal. *Behavioural Neuroscience* **120**: 93-102.
- KRUK, M. R., WESTPHAL, K.G., VAN ERP, A.M., VAN ASPEREN, J., CAVE, B.J., SLATER, E., DE KONING, J. and HALLER, J. (1998) The hypothalamus: cross-roads of endocrine and behavioural regulation in grooming and aggression. *Neuroscience and biobehavioural reviews* **23**: 163-177.
- KUCHEL, O. (1991) Stress and catecholamines. In: Stress Revisited. I. Neuroendocrinology of stress. Jasmin G. and M. Cantin eds. Methods achieve Exp Pathol. Basel Karger. Vol. **14**. pp 80-103.
- KUCHEL, O., BUU, N.T. and WANG, P.C. (1982) Stress, catecholamines and hypertension: Respective roles of free and conjugated catecholamines. *Stress* **3**: 22-31.
- KUDRIAVTSEVA, N.N., NIKULINA, E.M. and POPOVA, N.K. (1988) Participation of the dopamine of the striatum and the nucleus accumbens in the formation of aggressive and submissive behaviours in mice. *Zhurnal vysshej nervnoj deiatelnosti imeni I P Pavlova* **38**: 1168-1170.
- KUIKKA, J.T., TIHONEN, J., BERGSTROM, K.A., KARHU, J., RASANEN, P. and ERONEN, M. (1998) Abnormal structure of human striatal dopamine re-uptake sites in habitually violent alcoholic offenders: a fractal analysis. *Neuroscience Letter* **253**: 195-197.

- KULIKOV, A.V., ZHANAEVA, E.L.U. and POPOVA, N.K. (1989) Change in tryptophan hydroxylase activity in the brain of silver foxes and wild Norway rats in the course of selection according to behaviour. *Genetika* **25**: 346-350.
- LAMONT, S.J. (1994) Poultry immunogenetics: which way do we go? *Poultry Science* **73**: 1044-1048.
- LANG, G.F., ETCHES, R.J. and WALTON, J.S. (1984) Effects of luteinizing hormone, progesterone, testosterone, estradiol and corticosterone on ovulation and luteinizing hormone release in hens treated with aminoglutethimide. *Biology of Reproduction* **30**: 278-288.
- LAMPUGNANI, M.G., BUCZKO, W., CECI, A., MENNINI, A. and DE GAETANO, G. (1986) Normal serotonin uptake by blood platelets and brain synaptosomes but selective impairment of platelet serotonin storage in mice with Chediack-Higashi syndrome. *Life of Science* **38**: 2193-2198.
- LECHIN, F., VAN DER DIJS, B., OROZCO, B., LECHIN, A.E., BAEZ, S., LECHIN, M.E., BENAÏM, M., ACOSTA, E., AROCHA, L., JIMENEZ, V., LEON, G. and GARCIA, Z. (1996) Plasma neurotransmitters, blood pressure and heart rate during supine resting, orthostasis and moderate exercise in severely ill patients: a model of failing to cope with stress. *Psychotherapy and Psychosomatics* **65**: 129-136.
- LENDERS, J. W.M., BRUNER, H.G., MURPHY, D.L. and EISENHOFER, G. (1998) Genetic deficiencies of monoamine oxidase enzymes: A key to understanding the function of the enzymes in humans. In: Chen, K. and J.C. Shih. *Advances in Pharmacology*. Academic Press. Vol. **42**, pp. 297-301.
- LEWIS, M.H., GARIEPY, J.L., GENDREAU, P., NICHOLS, D.E. and MAILMAN, R.B. (1994) Social reactivity and D1 dopamine receptors: studies in mice selectively bred for high and low levels of aggression. *Neuropsychopharmacology* **10**: 115-122.
- LORRAIN, D.S., MATUSZEWICH, L. and HULL, E.M. (1998) 8-OH-DPAT influences extracellular levels of serotonin and dopamine in the medial preoptic area of male rats. *Brain Research* **790**: 217-223.
- MACNAMEE, M.C. and SHARP, P.J. (1989) The functional activity of hypothalamic dopamine in broody bantam hens. *The Journal of Endocrinology* **121**: 67-74.
- MARAZZITI, D., PALEGO, L., SILVESTRI, S., PRESTA, S., BALESTRI, C., BATISTINI, A. and CONTI, L. (1996) Platelet abnormalities in aggressive subjects with mental deficiency. *Journal of Psychiatry and Neuroscience* **21**: 109-113.
- MARTIN, F. and ARTIGAS, F. (1992) Simultaneous effects of p-chloroamphetamine, d-fenfluramine, and reserpine on free and stored 5-hydroxytryptamine in brain and blood. *Journal of Neurochemistry* **59**: 1138-44.
- MARTIN, W.H., ROGOL, A.D., KAISER, D.L. and THORNER, M.O. (1981) Dopaminergic mechanisms and luteinizing hormone (LH) secretion. II. Differential effects of dopamine and bromocriptine on LH release in normal women. *The Journal of Clinical Endocrinology and Metabolism* **52**: 650-656.
- MASWOOD, N., CALDAROLA-PASTUSZKA, M. and UPHOUSE, L. (1998) Functional integration among 5-hydroxytryptamine receptor families in the control of female rat sexual behaviour. *Brain Research* **802**: 98-103.
- MENCH, J.A. (1992) Introduction: applied ethology and poultry science. *Poultry Science* **71**: 631-633.
- MENCH, J.A. and SHEA-MOORE, M. (1995) Moods, minds and molecules: the neurochemistry of social behaviour. *Applied Animal Behaviour Science* **44**: 99-118.
- MENCH, J.A. and DUNCAN, I.J. (1998) Poultry welfare in North America: opportunities and challenges. *Poultry Science* **77**: 1763-1765.
- MENDELSON, S.D. (1992) A review and reevaluation of the role of serotonin in the modulation of lordosis behaviour in the female rat. *Neuroscience and Biobehavioural Reviews* **16**: 309-350.
- MENDELSON, S.D. and GORZALKA, B.B. (1985) A facilitatory role for serotonin in the sexual behaviour of the female rat. *Pharmacology, Biochemistry and Behaviour* **22**: 1025-1033.
- MENDELSON, S.D. and MCEWEN, B.S. (1992) Autoradiographic analyses of the effects of adrenalectomy and corticosterone on 5-HT1A and 5-HT1B receptors in the dorsal hippocampus and cortex of the rat. *Neuroendocrinology* **55**: 444-450.
- MERSMANN, H.J. (1998) Overview of the effects of beta-adrenergic receptor agonists on animal growth including mechanisms of action. *Journal of Animal Science* **76**: 160-172.
- MICZEK, K.A., WEERTS, E., HANEY, M. and TIDEY, J. (1994) Neurobiological mechanisms controlling aggression: preclinical developments for pharmacotherapeutic interventions. *Neuroscience and Biobehavioural Reviews* **18**: 97-110.
- MILLARD, S.A., COSTA, E. and GAL, E.M. (1972) On the control of brain serotonin turnover rate by end product inhibition. *Brain Research* **40**: 545-551.
- MILLS, A.D. and FAURE, J.M. (1991) Divergent selection for duration of tonic immobility and social reinstatement behaviour in Japanese quail (*Coturnix coturnix japonica*) chicks. *Journal of Comparative Psychology* **105**: 25-38.
- MOFFITT, T.E., BRAMMER, G.L., CASPI, A., FAWCETT, J.P., RALEIGH, M., YUWILER, A. and SILVA, P. (1998) Whole blood serotonin relates to violence in an epidemiological study. *Biological Psychiatry* **43**: 446-457.
- MORELLO, H., CALIGARIS, L., HAYMAL, B. and TALEISNIK, S. (1992) Daily variations in the sensitivity of proestrous LH surge in the inhibitory effect of intraventricular injection of 5-HT or GABA in rats. *Canadian Journal of Physiology and Pharmacology* **70**: 447-451.

- MUCK-SELER, D., PIVAC, N., JAKOVLJEVIC, M. and BRZOVIC, Z. (1999) Platelet serotonin, plasma cortisol, and dexamethasone suppression test in schizophrenic patients. *Biological Psychiatry* **45**: 1433-1439.
- MUIR, W.M. (1996) Group selection for adaptation to multiple-hen cages: selection program and direct responses. *Poultry Science* **75**: 447-458.
- MUIR, W.M. AND CRAIG, J.V. (1998) Improving animal well-being through genetic selection. *Poultry Science* **77**: 1781-1788.
- MUNCK, A., GUYRE, P.M. and HOLBROOK, N.J. (1984) Physiological functions of glucocorticoids in stress and their relation to pharmacological actions. *Endocrine Reviews* **5**: 25-44.
- NAGATSUKA, Y. (1983) The regulation of pituitary gonadotropin release of the frontal lobe neocortex. (II). In relation to serotonergic neurons. *Nippon Naibunpi Gakkai Zasshi* **59**: 1874-1883.
- NANKOVA, B.B. AND SABBAN, E.L. (1999) Multiple signalling pathways exist in the stress-triggered regulation of gene expression for catecholamine biosynthetic enzymes and several neuropeptides in the rat adrenal medulla. *Acta Physiologica Scandinavica* **167**: 1-9.
- NAUMENKO, E.V., POPOVA, N.K. and IVANOVA, L.N. (1987) Neuroendocrine and neurochemical mechanisms of the domestication of animals. *Genetika* **23**: 1011-1025.
- NAUMENKO, E.V., POPOVA, N.K., NIKULINA, E.M., DYGAŁO, N.N., SHISHKINA, G.T., BORODIN, P.M. and MARKEL, A.L. (1989) Behaviour, adrenocortical activity, and brain monoamines in Norway rats selected for reduced aggressiveness towards man. *Pharmacology, Biochemistry, and Behaviour* **33**: 85-91.
- NESTOR, K.E., ANDERSON, J.W. and PATTERSON, R.A. (2000) Genetics of growth and reproduction in the turkey. 14. Changes in genetic parameters over thirty generations of selection for increased body weight. *Poultry Science* **79**: 445-452.
- NIKULINA, E.M. and KAPRALOVA, N.S. (1991) The role of dopamine receptors in controlling mouse aggressivity: the genotype dependence. *Zhurnal vysshej nervnoj deiatelnosti imeni I P Pavlova* **41**: 734-740.
- NIKULINA, E.M. and KAPRALOVA, N.S. (1992) Role of dopamine receptors in the regulation of aggression in mice; relationship to genotype. *Neuroscience and Behavioural Physiology* **22**: 364-369.
- NOMURA, M. and NOMURA, Y. (2006) Psychological, neuroimaging, and biochemical studies on functional association between impulsive behaviour and the 5-HT_{2A} receptor gene polymorphism in humans. *Annals of the New York Academy Sciences* **1086**: 134-143.
- OLIVIER, B., VAN OORSCHOT, R. and WALDINGER, M.D. (1998) Serotonin, serotonergic receptors, selective serotonin reuptake inhibitors and sexual behaviour. *International Clinical Psychopharmacology* **13** (Suppl 6): S9-14.
- OQUENDO, M.A. and MANN, J.J. (2000) The biology of impulsivity and suicidal. *The Psychiatric Clinics of North America* **23**: 11-25.
- OSSOWSKA, G., KLENK-MAJEWSKA, B. and ZEBROWSKA-LUPINA, I. (1996) Acute effect of dopamine agonists and some antidepressants in stress-induced deficit of fighting behaviour. *Polish Journal of Pharmacology* **48**: 403-408.
- OWENS, R.E., FLEEGER, J.L. and HARMS, P.G. (1980) Evidence for central nervous system (CNS) involvement in inhibition of luteinizing hormone (LH) release by dopamine receptor stimulation. *Endocrine Research Communications* **7**: 99-105.
- PANT, K.K. and NATH, C. (1993) Dopaminergic involvement in the effects of piracetam on foot shock induced aggression in mice. *The Indian Journal of Medical Research* **98**: 155-159.
- PARMIGIANI, S., PALANZA, P., ROGERS, J. and FERRARI, P.F. (1999) Selection, evolution of behaviour and animal models in behavioural neuroscience. *Neuroscience and Biobehavioural Reviews* **23**: 957-969.
- PAUL, S.M., REHAVI, M., SKOLNICK, P. and GOODWIN, F.K. (1980) Demonstration of specific "high affinity" binding sites for [3H] imipramine on human platelets. *Life Sciences* **26**: 953-959.
- PIERCECCHI-MARTI, M.D., LEONETTI, G., PELISSIER, A.L., CONRATH, J., CIANFARANI, F. and VALLI, M. (1999) Evaluation of biological stress markers in police officers. *Medicine and Law* **18**: 125-144.
- PIETRASZEK, M.H., TAKADA, Y., YAN, D., URANO, T., SERIZAWA, K. and TAKADA, A. (1992) Relationship between serotonergic measures in periphery and the brain of mouse. *Life Sciences* **51**: 75-82.
- PLETSCHER, A. (1988) Platelets as models: use and limitations. *Experientia* **44**: 152-155.
- POLLARD, T.M. (1997) Physiological consequences of everyday psychosocial stress. *Collegium Antropologicum* **21**: 17-28.
- POPOVA, N.K. (2006) From genes to aggressive behaviour: the role of serotonergic system. *Bioessays* **28**: 495-503.
- POPOVA, N.K., VOITENKO, N.N. and TRUT, L.N. (1975) Changes in the content of serotonin and 5-hydroxyindoleacetic acid in the brain during selection of silver foxes according to behaviour. *Doklady Akademii nauk SSSR* **223**: 1498-1500.
- POPOVA, N.K., KUDRIAVTSEVA, N.N. and LUBSANOVA, S.D. (1978) Genetic control of the tissue concentration of serotonin in mice. *Genetika* **14**: 1804-1808.
- POPOVA, N.K., KULIKOV, A.V., AVGUSTINOVICH, D.F., VOITENKO, N.N. and TRUT, L.N. (1997) Effect of domestication of the silver fox on the main enzymes of serotonin metabolism and serotonin receptors. *Genetika* **33**: 370-374.

- POPOVA, N.K., BARYKINA, N.N., PLIUSNINA, I.Z., ALEKHINA, T.A. and KOLPAKOV, V.G.** (1999) Manifestation of fear response in rats genetically predisposed to various kinds of defense behaviour. *Rossiĭskii fiziologicheskii zhurnal imeni I.M. Sechenova / Rossiĭskaya akademiya nauk* **85**: 99-104.
- RALEIGH, M.J., MCGUIRE, M.T., BRAMMER, G.L., POLLACK, D.B. and YUWILER, A.** (1991) Serotonergic mechanisms promote dominance acquisition in adult male vervet monkeys. *Brain Research* **559**: 181-190.
- RENDEN, J.A., LIEN, R.J., OATES, S.S. and BILGILI, S.F.** (1994) Plasma concentrations of corticosterone and thyroid hormones in broilers provided various lighting schedules. *Poultry Science* **73**: 186-193.
- RICHFIELD, E.K., YOUNG, A.B. and PENNEY, J.B.** (1987) Comparative distribution of dopamine D-1 and D-2 receptors in the basal ganglia of turtles, pigeons, rats, cats, and monkeys. *Journal of Comparative Neurology* **262**: 446-463.
- ROSENBURG, T.B., TUYTTENS, F.A., SONCK, B., DE REU, K., HERMAN, L. and ZOONS, J.** (2005) Welfare, health, and hygiene of laying hens housed in furnished cages and in alternative housing systems. *Journal Applied Animal Welfare Science* **8**: 211-226.
- ROLIH, C.A. and OBER, K.P.** (1995) The endocrine response to critical illness. *The Medical Clinics of North America* **79**: 211-224.
- ROTS, N.Y., COOLS, A.R., DE JONG, J. and DE KLOET, E.R.** (1995) Corticosteroid feedback resistance in rats genetically selected for increased dopamine responsiveness. *Journal of Neuroendocrinology* **7**: 153-161.
- SAPOLSK, R.M.** (1990) Stress in the wild. *Scientific American* **262**: 116-123.
- SAPOLSKY, R.M. and MEANEY, M.J.** (1986) Maturation of the adrenocortical stress response: neuroendocrine control mechanisms and the stress hyporesponsive period. *Brain Research* **396**: 64-76.
- SAVORY C.J. and MANN, J.S.** (1997) Is there a role for corticosterone in expression of abnormal behaviour in restricted-fed fowls? *Physiology and Behaviour* **62**: 7-13.
- SCOTT, P.A., CIERPIAL, M.A., KILTS, C.D. and WEISS, J.M.** (1996) Susceptibility and resistance of rats to stress-induced decreases in swim-test activity: a selective breeding study. *Brain Research* **725**: 217-230.
- SERVO, L.I. and NAUMANKO, E.V.** (1990) The brain catecholamine systems in the regulation of dominance. *Zhurnal vyssheĭ nervnoĭ deiatelnosti imeni I P Pavlova* **40**: 490-496.
- SGOIFO, A., DE BOER, S.F., HALLER, J. and KOOLHAAS, J.M.** (1996) Individual differences in plasma catecholamine and corticosterone stress responses of wild-type rats: relationship with aggression. *Physiology and Behaviour* **60**: 1403-1407.
- SHAIKH, M.B., LU, C.L., MACGREGOR, M. and SIEGEL, A.** (1991) Dopaminergic regulation of quiet biting attack behaviour in the cat. *Brain Research Bulletin* **27**: 725-730.
- SHARP, P.J. and BEUVING, G.** (1978) The role of corticosterone in the ovulatory cycle of the hen. *Journal of Endocrinology* **78**: 195-200.
- SHARP, P.J., MACNAMEE, M.C., TALBOT, R.T., STERLING, R.J. and HALL, T.R.** (1984) Aspects of the neuroendocrine control of ovulation and broodiness in the domestic hen. *Journal of Experimental Zoology* **232**: 475-483.
- SHIVELY, C. and KAPLAN, J.** (1984) Effects of social factors on adrenal weight and related physiology of *Macaca fascicularis*. *Physiology and Behaviour* **33**: 777-782.
- SIEGEL, H.S.** (1971) Cold-induced necrosis in toes of chicks receiving daily dosages of adrenocorticotropin. *General and Comparative Endocrinology* **16**: 281-287.
- SIEGEL, A., ROELING, T.A., GREGG, T.R. and KRUK, M.R.** (1999) Neuropharmacology of brain-stimulation-evoked aggression. *Neuroscience and Biobehavioural Reviews* **23**: 359-389.
- SIEGEL, H.S.** (1995) Gordon Memorial Lecture. Stress, strains and resistance. *Bristol Poultry Science* **36**: 3-22.
- SIEGEL, P.B. and DUNNINGTON, E.A.** (1997) Genetic selection strategies—population genetics. *Poultry Science* **76**: 1062-1065.
- SIROTKIN, A.V. and SCHAEFFER, H.J.** (1997) Direct regulation of mammalian reproductive organs by serotonin and melatonin. *Journal of Endocrinology* **154**: 1-5.
- SNIDER, S.R. and KUCHEL, O.** (1983) Dopamine: an important neurohormone of the sympathoadrenal system: Significance of increased peripheral dopamine release for the human stress response and hypertension. *Endocrine Reviews* **4**: 291-309.
- SOTOWSKA-BROCHOCKA, J., MARTYNSKA, L. and LICHT, P.** (1994) Dopaminergic inhibition of gonadotropic release in hibernating frogs, *Rana temporaria*. *General and Comparative Endocrinology* **93**: 192-196.
- STAHL, S.M.** (1977) The human platelet. A diagnostic and research tool for the study of biogenic amines in psychiatric and neurologic disorders. *Archives of General Psychiatry* **34**: 509-516.
- STAHL, S.M., CIARANELL, R.D. and BERGER, P.A.** (1982) Platelet serotonin in schizophrenia and depression. In: Ho, B. T., Schoolar, J.C. and Usdin, E., eds. *Serotonin in biological psychiatry*. New York: Raven Press.
- STANULIS, E.D., MATULKA, R.A., JORDAN, S.D., ROSECRANS, J.A. and HOLSAPPLE, M.P.** (1997) Role of corticosterone in the enhancement of the antibody response after acute cocaine administration.

- Journal of Pharmacology and Experimental Therapeutics* **280**: 284-291.
- STEKELIS, H.D., RALEIGH, M.J., KLING, A.S. and TACHIKI, K.** (1986) Biochemical and hormonal correlates of dominance and social behaviour in all-male groups of squirrel monkeys (*Saimiri sciureus*). *American Journal of Primatology* **11**: 133-145.
- SUMMERS, C.H., LARSON, E.T., RONAN, P.J., HOFMANN, P.M., EMERSON, A.J. and RENNER, K.J.** (2000) Serotonergic responses to corticosterone and testosterone in the limbic system. *General and Comparative Endocrinology* **117**: 151-159.
- TABLANTE, N.L., VAILLANCOURT, J.P., MARTIN, S.W., SHOUKRI, M. and ESTEVEZ, I.** (2000) Spatial distribution of cannibalism mortality in commercial laying hens. *Poultry Science* **79**: 705-708.
- TAKADA, A., URANO, T., YOSHIDA, M. and TAKADA, Y.** (1996) Comparison of changes in serotonergic measures in whole blood or plasma and brain in rats given nicotine and/or stresses. *Polish Journal of Pharmacology* **48**: 173-177.
- TASAKA, K., MIYAKE, A., SAKUMOTO, T. and AONO, T.** (1985) Dopamine decreases release of luteinizing hormone releasing hormone from superfused rat mediobasal hypothalamus. *Journal of Endocrinological Investigation* **8**: 373-376.
- TECOTT, L.H. and BARONDES, S.H.** (1996) Genes and aggressiveness: Behavioural genetics. *Current Biology* **6**: 238-240.
- TEMPEL, D.L. and LEIBOWITZ, S.F.** (1994) Adrenal steroid receptors: interactions with brain neuropeptide systems in relation to nutrient intake and metabolism. *Journal of Neuroendocrinology* **6**: 479-501.
- THIERY, J.C., GAYRARD, V., LE CORRE, S., VIGUIE, C., MARTIN, G.B., CHEMINEAU, P. and MALPAUX, B.** (1995) Dopaminergic control of LH secretion by the A15 nucleus in anoestrous ewes. *Journal of Reproduction and Fertility Suppl.* **49**: 285-296.
- THOMPSON, D.L., ELGERT, K.D., GROSS, W.B. and SIEGEL, P.B.** (1980) Cell-mediated immunity in Marek's disease virus-infected chickens genetically selected for high and low concentrations of plasma corticosterone. *American Journal of Veterinary Research* **41**: 91-96.
- TUOMISTO, J. and MANNISTO, P.** (1985) Neurotransmitter regulation of anterior pituitary hormones. *Pharmacological Reviews* **37**: 249-332.
- UNIS, A.S., COOK, E.H., VINCENT, J.G., GJERDE, D.K., PERRY, B.D., MASON, C. and MITCHELL, J.** (1997) Platelet serotonin measures in adolescents with conduct disorder. *Biological Psychiatry* **42**: 553-559.
- VADASZ, C., KOBOR, G. and LAJTHA, A.** (1992) Motor activity and the mesotelencephalic dopamine function. I. High-resolution temporal and genetic analysis of open-field behaviour. *Behavioural Brain Research* **48**: 29-39.
- VAN ERP, A.M. and MICZEK, K.A.** (2006) Increased accumbal dopamine during daily alcohol consumption and subsequent aggressive behaviour in rats. *Psychopharmacology* (in press).
- VALZELLI, L.** (1984) Reflections on experimental and human pathology of aggression. *Prog Neuropsychopharmacol Biological Psychiatry* **8**: 311-325.
- VIRKKUNEN, M. and LINNOILA, M.** (1997) Serotonin in early-onset alcoholism. *Recent Developments in Alcoholism* **13**: 173-189.
- VOGEL, W.H. and HARRIS, N.** (1991) Learning and memory of a water T-maze by rats selectively bred for low or high plasma catecholamine stress responses. *Behavioural Neural Biology* **56**: 113-117.
- VOLAVKA, J., BILDER, R. and NOLAN, K.** (2004) Catecholamines and aggression: the role of COMT and MAO polymorphisms. *Annals of the New York Academy Sciences* **1036**: 393-398.
- VOLLMER, R.R.** (1996) Selective neural regulation of epinephrine and norepinephrine cells in the adrenal medulla — cardiovascular implications. *Clinical and Experimental Hypertension* **18**: 731-751.
- WALKER, E.A., YAMAMOTO, T., HOLLINGSWORTH, P.J., SMITH, C.B. and WOODS, J.H.** (1991) Discriminative-stimulus effects of quipazine and 1-5-hydroxytryptophan in relation to serotonin binding sites in the pigeon. *Journal of Pharmacology Experimental Therapeutics* **259**: 772-782.
- WATT, M.J., FORSTER, G.L., KORZAN, W.J., RENNER, K.J. and SUMMERS, C.H.** (2006) Rapid neuroendocrine responses evoked at the onset of social challenge. *Physiology and Behaviour* (in press).
- WEIGER, W.A.** (1997) Erotonergic modulation of behaviour: a phylogenetic overview. *Biological Reviews of the Cambridge Philosophical Society* **72**: 61-95.
- WESTERGAARD, G.C., SUOMI, S.J., HIGLEY, J.D. and MEHLMAN, P.T.** (1999) CSF 5-HIAA and aggression in female macaque monkeys: species and interindividual differences. *Psychopharmacology* **146**: 440-446.
- WILSON, C.A. and HUNTER, A.J.** (1985) Progesterone stimulates sexual behaviour in female rats by increasing 5-HT activity on 5-HT₂ receptors. *Brain Research* **333**: 223-229.
- WOLF, A., CALDAROLA-PASTUSZKA, M. and UPHOUSE, L.** (1998) Facilitation of female rat lordosis behaviour by hypothalamic infusion of 5-HT(2A/2C) receptor agonists. *Brain Research* **779**: 84-95.
- WRASE, J., REIMOLD, M., PULS, I., KIENAST, T. and HEINZ, A.** (2006) Serotonergic dysfunction: brain imaging and behavioural correlates. *Cognitive Affective and Behavioural Neuroscience* **6**: 53-61.
- YAN, D., URANO, T., PIETRASZEK, M.H., SHIMOYAMA, I., UEMURA, K., KOJIMA, Y., SAKAKIBARA, K., SERIZAWA, K., TAKADA, Y. and TAKADA, A.** (1993) Correlation between serotonergic measures in cerebrospinal fluid and blood of subhuman primate. *Life of Science* **52**: 745-749.

YOUNG, W.F. JR, LAWS E.R. JR, SHARBROUGH, F.W. and WEINSHILBOUM, R.M. (1986) Human monoamine oxidase. Lack of brain and platelet correlation. *Archives of General Psychiatry* **43**: 604-609.

Table 1 Effect of genetic selection on survivability and productivity in hens.

Group	Survivability		Productivity		
	Longevity (Days)	Mortality (%)	Egg Number (Per Cage)	Egg Mass (Per Cage, g/day)	Egg Weight (g/each)
KGB	363 ^a ± 0.4	1.3 ^a ± 0.1	3543 ^a ± 139	574 ^a ± 24	59.4 ^a ± 0.6
LGPS	193 ^b ± 21.	8.6 ^b ± 0.5	1297 ^b ± 140	208 ^b ± 22	58.9 ^a ± 0.8
Overall	329 ^c ± 2.0	2.3 ^c ± 0.1	2850 ^c ± 26.9	450 ^c ± 4.4	57.8 ^a ± 0.1

1) ^{a,b,c}Lines with different superscript are significantly different ($P < 0.01$).

2) Values represent the Mean ± SEM. The line "Overall" was based on the mean of 768 colony cages (9216 birds), and the KGB and LGP line were based on the mean of 12 cages (144 birds) of each line, respectively.

Table 2 Genetic selection-induced alterations in blood concentrations of serotonin catecholamines, and corticosterone in hens.

Groups	DA (ng/ml)	EP (ng/ml)	NE (ng/ml)	EP/NE (Ratio, %)	CORT (mg/ml)
KGB	0.59 ± 0.08 ^a	0.30 ± 0.06 ^a	0.86 ± 0.12	34.0 ^a	1.87 ± 0.19
LGPS	2.42 ± 0.76 ^b	0.59 ± 0.13 ^b	0.84 ± 0.13	72.5 ^b	1.49 ± 0.21

^{a,b}Means within a column with different superscript are statistically different ($P < 0.01$).

The KGB and LGP lines were selected from high and low productivity and survivability resulting from cannibalism and flightiness. 5-HT, serotonin; DA, dopamine; EP, Epinephrine; and NE, Norepinephrine.




Figure 1 Quantity analysis of blood serotonin levels using HPLC. a) calibration curve, and b) effect of genetic selection on the blood concentration of serotonin. Asterisk indicates significantly different between selected lines. $P < 0.01$.




Figure 2 Examples of effects of genetic selection on blood concentrations of dopamine, epinephrine and norepinephrine. a) calibration curve, b) data from the KGB bird, and c) data from the LGP bird. The peak height of each neurochemistry in each graph was referred to its concentration based on the calibration cure, respectively.